

$B \rightarrow K\pi$ decays

T. N. Pham

*Centre de Physique Théorique, CNRS
Ecole Polytechnique, 91128 Palaiseau, Cedex, France*
(Dated: August 19, 2009)

Though QCD Factorization(QCDF) could produce sufficiently large $B \rightarrow K\pi$ branching ratios close to experiments, the predicted direct CP asymmetry in $\bar{B}^0 \rightarrow K^+\pi^-$ decay is however still quite below experiment, even with a large negative phase in the annihilation term, according to existing calculations. This suggests the presence of an additional strong phase in the decay amplitude which could come from the long-distance final state interactions(FSI) like charming penguin. In this paper, we show that, by adding to the $B \rightarrow K\pi$ decay QCDF amplitude, a real part and an absorptive part with a strength 10% and 30% of the penguin amplitude, respectively, we could bring the $B \rightarrow K\pi$ branching ratios and the $\bar{B}^0 \rightarrow K^-\pi^+$ CP asymmetry close to the measured values. We also find that the color-allowed electroweak penguin is appreciable and that from the QCDF electroweak-strong penguin interference terms and the measured $B^- \rightarrow \bar{K}^0\pi^-$ we obtain $(9.0 \pm 0.3) \times 10^{-6}$ for the $\bar{B}^0 \rightarrow \bar{K}^0\pi^0$ branching ratio, a bit lower than experiment. Similarly, the predicted value for $B^- \rightarrow K^-\pi^0$ in terms of the $\bar{B}^0 \rightarrow K^-\pi^+$ branching ratio agrees well with experiment. This suggests that the measured value for $\bar{B}^0 \rightarrow \bar{K}^0\pi^0$ in fact should be lower than the current measured value.

PACS numbers: 13.25Hw, 12.38.Bx

I. INTRODUCTION

Existing calculations of the penguin-dominated $B \rightarrow K\pi$ decays in QCD Factorization(QCDF) [1, 2, 3] seem to produce $B \rightarrow K\pi$ branching ratios more or less in agreement with experiments, with a moderate annihilation contribution, but the predicted direct CP asymmetry in $\bar{B}^0 \rightarrow K^-\pi^+$ decay is smaller than experiment by a factor of 2 or more, even with a large negative strong phase for the annihilation terms as in scenario S4 [2]. In fact in QCDF, and also in naive Factorization model [4], the absorptive part(the imaginary part) of the short-distance Wilson coefficients in the non leptonic decay effective Hamiltonian are negative and of the same sign as the real part, the direct CP asymmetry for $\bar{B}^0 \rightarrow \pi^+\pi^-$ and $\bar{B}^0 \rightarrow K^-\pi^+$ are negative and positive, respectively, in opposite sign with measurements[5]. To reverse the sign of the CP asymmetry and to produce a large CP asymmetry, one would need a large annihilation term with a large negative phase. Another possibility is a long-distance final state interaction(FSI) term, probably mainly absorptive, to generate a large strong phase, but not to overestimate the branching ratios, like charming penguin due to $B \rightarrow D_s D \rightarrow K\pi$ and $B \rightarrow D_s^* D^* \rightarrow K\pi$ inelastic FSI effect [6, 7, 8, 9, 10, 11] which are CKM-favored, color-allowed processes and could produce a large absorptive part in $B \rightarrow K\pi$ decays. In this paper, we show that by adding to the QCDF amplitude, a penguin-like amplitude with a real and absorptive part with a strength 10% and 30%, respectively, of the penguin amplitude, we could obtain the $B \rightarrow K\pi$ branching ratios and the $\bar{B}^0 \rightarrow K^-\pi^+$ CP asymmetry close to experiments. We also find that the color-allowed electroweak penguin(EW) from the operators O_7 and O_9 contribute appreciably through the strong penguin-EW penguin in-

terference terms in $B^- \rightarrow K^-\pi^0$ and $\bar{B}^0 \rightarrow \bar{K}^0\pi^0$ decays. Then using only the QCDF expressions for the color-allowed tree-penguin(T-P), color-suppressed tree-penguin(C-P), color-allowed electroweak-penguin(EW-P) interference terms and the measured $\mathcal{B}(B^- \rightarrow \bar{K}^0\pi^-)$ and $\mathcal{B}(\bar{B}^0 \rightarrow K^-\pi^+)$, we could predict $\mathcal{B}(B^- \rightarrow K^-\pi^0)$ and $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$. We find that the predicted $\mathcal{B}(B^- \rightarrow K^-\pi^0)$ agrees well with experiment while the predicted $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$ is below the measured value. This strongly suggests that the measured $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$ is a bit too high as shown in the following. In the next section we will present numerical results for the $B \rightarrow K\pi$ branching ratios and CP asymmetry in QCDF factorization with the addition of a small penguin-like amplitude to generate a CP asymmetry for $\bar{B}^0 \rightarrow K^-\pi^+$ decay. We then give prediction for $\mathcal{B}(B^- \rightarrow K^-\pi^0)$ and $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$ from the electroweak penguin and strong penguin interference terms.

II. $B \rightarrow K\pi$ DECAY IN QCD FACTORIZATION

The $B \rightarrow M_1 M_2$ decay amplitude in QCDF is given by [1, 2]:

$$\mathcal{A}(B \rightarrow M_1 M_2) = \frac{G_F}{\sqrt{2}} \sum_{p=u,c} V_{pb} V_{ps}^* \times \left(- \sum_{i=1}^{10} a_i^p \langle M_1 M_2 | O_i | B \rangle_H + \sum_i^{10} f_B f_{M_1} f_{M_2} b_i \right), \quad (1)$$

where the QCD coefficients a_i^p contain the vertex corrections, penguin corrections, and hard spectator scattering contributions, the hadronic matrix elements $\langle M_1 M_2 | O_i | B \rangle_H$ of the tree and penguin operators O_i are given by factorization model [4, 12], b_i are annihilation

contributions. The values for $a_i^p, p = u, c$, computed from the expressions in [1, 2] at the renormalization scale $\mu = m_b$, with $m_b = 4.2 \text{ GeV}$, are :

$$\begin{aligned} a_4^c &= -0.033 - 0.013i + 0.0009 \rho_H \exp(i\phi_H), \\ a_4^u &= -0.027 - 0.017i + 0.0009 \rho_H \exp(i\phi_H), \\ a_6^c &= -0.045 - 0.003i, \quad a_6^u = -0.042 - 0.013i, \\ a_8^c &= -0.0004 - 0.0001i, \quad a_8^u = 0.0004 - 0.0001i, \\ a_{10}^c &= -0.0011 - 0.0001i - 0.0006 \rho_H \exp(i\phi_H), \\ a_{10}^u &= -0.0011 + 0.0006i - 0.0006 \rho_H \exp(i\phi_H). \end{aligned} \quad (2)$$

for $i = 4, 6, 8, 10$. For other coefficients, $a_i^u = a_i^p = a_i$:

$$\begin{aligned} a_1 &= 1.02 + 0.015i - 0.012 \rho_H \exp(i\phi_H), \\ a_2 &= 0.156 - 0.089i + 0.074 \rho_H \exp(i\phi_H), \\ a_3 &= 0.0025 + 0.0030i - 0.0024 \rho_H \exp(i\phi_H), \\ a_5 &= -0.0016 - 0.0034i + 0.0029 \rho_H \exp(i\phi_H), \\ a_7 &= -0.00003 - 0.00004i - 0.00003 \rho_H \exp(i\phi_H) \\ a_9 &= -0.009 - 0.0001i + 0.0001 \rho_H \exp(i\phi_H). \end{aligned} \quad (3)$$

where the complex parameter $\rho_H \exp(i\phi_H)$ represents the end-point singularity contribution in the hard-scattering corrections $X_H = (1 + \rho_H \exp(i\phi_H)) \ln(\frac{m_B}{\Lambda_h})$ [1, 2].

For the annihilation terms, we have :

$$\begin{aligned} b_2 &= -0.0041 - 0.0071 \rho_A \exp(i\phi_A) - 0.0019 (\rho_A \exp(i\phi_A))^2, \\ b_3 &= -0.0071 - 0.016 \rho_A \exp(i\phi_A) - 0.0093 (\rho_A \exp(i\phi_A))^2, \\ b_3^{ew} &= -0.00012 - 0.00016 \rho_A \exp(i\phi_A) \\ &\quad + 0.000003 (\rho_A \exp(i\phi_A))^2. \end{aligned} \quad (4)$$

where b_i are evaluated with the factor $f_B f_{M_1} f_{M_2}$ included and normalized relative to the factor $f_K F_0^{B\pi}(m_B^2 - m_\pi^2)$ in the factorisable terms, and ρ_A , like ρ_H , appears in the divergent annihilation term $X_A = (1 + \rho_A \exp(i\phi_A)) \ln(\frac{m_B}{\Lambda_h})$.

The $B \rightarrow K\pi$ decay amplitude with the factorisable part [4] and the annihilation term [1, 2, 13] are:

$$\begin{aligned} A(B^- \rightarrow K^- \pi^0) &= -i \frac{G_F}{2} f_K F_0^{B\pi}(m_K^2)(m_B^2 - m_\pi^2) \\ &\quad (V_{ub}V_{us}^* a_1 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*)[a_4 + a_{10} + (a_6 + a_8)r_\chi]) \\ &\quad -i \frac{G_F}{2} f_\pi F_0^{BK}(m_\pi^2)(m_B^2 - m_K^2) \\ &\quad \times \left(V_{ub}V_{us}^* a_2 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times \frac{3}{2}(a_9 - a_7) \right) \\ &\quad -i \frac{G_F}{2} f_B f_K f_\pi \\ &\quad \times [V_{ub}V_{us}^* b_2 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times (b_3 + b_3^{ew})] \end{aligned} \quad (5)$$

$$\begin{aligned} A(B^- \rightarrow \bar{K}^0 \pi^-) &= -i \frac{G_F}{\sqrt{2}} f_K F_0^{B\pi}(m_K^2)(m_B^2 - m_\pi^2) \\ &\quad + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \left[a_4 - \frac{1}{2}a_{10} + (a_6 - \frac{1}{2}a_8)r_\chi \right] \\ &\quad -i \frac{G_F}{\sqrt{2}} f_B f_K f_\pi \\ &\quad \times [V_{ub}V_{us}^* b_2 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times (b_3 + b_3^{ew})] \end{aligned} \quad (6)$$

and for $\bar{B}^0$:

$$\begin{aligned} A(\bar{B}^0 \rightarrow K^- \pi^+) &= -i \frac{G_F}{\sqrt{2}} f_K F_0^{B\pi}(m_K^2)(m_B^2 - m_\pi^2) \\ &\quad \left(V_{ub}V_{us}^* a_1 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*)[a_4 + a_{10} + (a_6 + a_8)r_\chi] \right) \\ &\quad -i \frac{G_F}{\sqrt{2}} f_B f_K f_\pi \left[(V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times (b_3 - \frac{b_3^{ew}}{2}) \right] \end{aligned} \quad (7)$$

$$\begin{aligned} A(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0) &= i \frac{G_F}{2} f_K F_0^{B\pi}(m_K^2)(m_B^2 - m_\pi^2) \\ &\quad \times (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \left[a_4 - \frac{1}{2}a_{10} + (a_6 - \frac{1}{2}a_8)r_\chi \right] \\ &\quad -i \frac{G_F}{2} f_\pi F_0^{BK}(m_\pi^2)(m_B^2 - m_K^2) \\ &\quad \left(V_{ub}V_{us}^* a_2 + (V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times \frac{3}{2}(a_9 - a_7) \right) \\ &\quad +i \frac{G_F}{2} f_B f_K f_\pi \left[(V_{ub}V_{us}^* + V_{cb}V_{cs}^*) \times (b_3 - \frac{b_3^{ew}}{2}) \right] \end{aligned} \quad (8)$$

where $r_\chi = \frac{2m_K^2}{(m_b - m_d)(m_d + m_s)}$ is the chirally-enhanced terms in the penguin O_6 matrix element. We also need the $B^- \rightarrow \pi^- \pi^0$ amplitude:

$$\begin{aligned} A(B^- \rightarrow \pi^- \pi^0) &= -i \frac{G_F}{2} f_\pi F_0^{B\pi}(m_\pi^2)(m_B^2 - m_\pi^2) \\ &\quad \left(V_{ub}V_{ud}^* (a_1 + a_2) + (V_{ub}V_{ud}^* + V_{cb}V_{cd}^*) \right. \\ &\quad \left. \times \frac{3}{2}(a_9 - a_7 + a_{10} + a_8 r_\chi) \right) \end{aligned} \quad (9)$$

We see that the $B \rightarrow K\pi$ decay amplitudes consist of a QCD penguin(P) $a_4 + a_6 r_\chi$, a color-allowed tree(T) a_1 , a color-suppressed tree(C) a_2 , a color-allowed electroweak penguin(EW) $a_9 - a_7$, a color-suppressed electroweak penguin(EWC) $a_{10} + a_8 r_\chi$ terms. Because of the relative large Wilson coefficients, the QCD penguin, the color-allowed tree and the color-allowed electroweak contributions give the major contribution in $B \rightarrow K\pi$ decays.

For the CKM matrix elements, since the inclusive and exclusive data on $|V_{ub}|$ differ by a large amount and the higher inclusive data exceeds the unitarity limit for $R_b = |V_{ud}V_{ub}^*|/|V_{cd}V_{cb}^*|$ with the current value $\sin(2\beta) = 0.681 \pm 0.025$ [14], we shall determine $|V_{ub}|$ from the more precise $|V_{cb}|$ data. We have [15]:

$$|V_{ub}| = \frac{|V_{cb}V_{cd}^*|}{|V_{ud}^*|} |\sin \beta| \sqrt{1 + \frac{\cos^2 \alpha}{\sin^2 \alpha}}. \quad (10)$$

With $\alpha = (99_{-9}^{+13})^\circ$ [14] and $|V_{cb}| = (41.78 \pm 0.30 \pm 0.08) \times 10^{-3}$ [16], we find

$$|V_{ub}| = 3.60 \times 10^{-3}. \quad (11)$$

in good agreement with the exclusive data in the range $|V_{ub}| = 3.33 - 3.51$ [16]. A recent UT fit also gives

$|V_{ub}| = (3.60 \pm 0.12) \times 10^{-3}$ [17]. The measurements of the $B_s - \bar{B}_s$ mixing also allow the extraction of $|V_{td}|$ from $B_d - \bar{B}_d$ mixing data. The current determination [18] gives $|V_{td}/V_{ts}| = (0.208_{-0.006}^{+0.008})$ which in turn can be used to determine the angle γ from the unitarity relation [15]:

$$|V_{td}| = \frac{|V_{cb}V_{cd}^*|}{|V_{tb}|} |\sin \gamma \sqrt{1 + \frac{\cos^2 \alpha}{\sin^2 \alpha}}|. \quad (12)$$

with $|V_{tb}| = 1$, we find $\gamma = 66^\circ$ which implies an angle $\alpha = 91.8^\circ$, in good agreement with the value found in the current UT-fit value of $(88 \pm 16)^\circ$ [19]. In the following in our $B \rightarrow K\pi$ decay calculations, we shall use the unitarity triangle values for $|V_{ub}|$ and γ . For other hadronic parameters we use the values in Table 1 of [2] and take $m_s(2\text{ GeV}) = 80\text{ MeV}$. For the $B \rightarrow \pi$ and $B \rightarrow K$ transition form factor, we use the current light-cone sum rules central value [20] :

$$F_0^{B\pi}(0) = 0.258, \quad F_0^{BK}(0) = 0.33 \quad (13)$$

The computed branching ratios and direct CP asymmetries, with $\rho_H = 1, \phi_H = 0$ and $\phi_A = -55^\circ$ as in scenario S4 of [2] are shown in Fig.1 and Fig.2 as function of ρ_A . For convenience we also give in Table I and Table II the computed values at $\rho_A = 1$ as in S4 with and without the additional penguin-like contribution δP .

Modes	$\delta P = 0$	$\delta P \neq 0$	Exp [21]
$B^- \rightarrow \pi^- \pi^0$	5.7	5.7	5.59 ± 0.4
$B^- \rightarrow K^- \pi^0$	10.3	12.7	12.9 ± 0.6
$B^- \rightarrow \bar{K}^0 \pi^-$	18.1	22.7	23.1 ± 1.0
$\bar{B}^0 \rightarrow K^- \pi^+$	15.5	19.7	19.4 ± 0.6
$\bar{B}^0 \rightarrow \bar{K}^0 \pi^0$	6.8	8.8	9.8 ± 0.6

TABLE I: The CP-averaged $B \rightarrow K\pi$ Branching ratios in unit of 10^{-6} in QCDF with and without additional penguin-like contribution δP and with $\rho_A = 1.0, \phi_A = -55^\circ$

Modes	$\delta P = 0$	$\delta P \neq 0$	Exp [21]
$B^- \rightarrow \pi^- \pi^0$	0.0	0.0	0.06 ± 0.05
$B^- \rightarrow K^- \pi^0$	0.02	-0.06	0.05 ± 0.025
$B^- \rightarrow \bar{K}^0 \pi^-$	0.004	0.01	-0.009 ± 0.025
$\bar{B}^0 \rightarrow K^- \pi^+$	-0.01	-0.08	$-0.098_{-0.010}^{+0.012}$
$\bar{B}^0 \rightarrow \bar{K}^0 \pi^0$	-0.02	0.01	-0.01 ± 0.010

TABLE II: The direct $B \rightarrow K\pi$ CP asymmetries in QCDF with and without additional penguin-like contribution δP and with $\rho_A = 1.0, \phi_A = -55^\circ$

As can be seen, without the additional penguin contribution, the computed branching ratios are slightly below experiments, but the CP asymmetries for $\bar{B}^0 \rightarrow K^- \pi^+$

is much smaller than the measured values. Considering various theoretical uncertainties, one could say that the real part of the $B \rightarrow K\pi$ decay amplitudes is more or less well described by QCDF, but the absorptive part of the amplitude needs some additional contribution to account for the observed CP asymmetries. As mentioned earlier, the factorisable contribution in the QCDF amplitude give the wrong sign for the CP asymmetries in $\bar{B}^0 \rightarrow K^- \pi^+$, one needs a sizeable absorptive part to reverse the sign of the asymmetries. As shown in Fig.1 and Fig.2, we would need an annihilation term with a very large ρ_A to produce correct $B \rightarrow K\pi$ branching ratios and $\bar{B}^0 \rightarrow K^- \pi^+$ asymmetry. This leaves us with charming penguin, as mentioned earlier, as a possible contribution to the large strong phase in $B \rightarrow K\pi$ decays. This possibility is considered in this paper by adding to the QCD penguin a small term:

$$\delta P = (ra_4 + ra_6 r_\chi)(d_1 + id_2) \quad (14)$$

where ra_4 and ra_6 are the real part of a_4 and a_6 , respectively. We note a previous fit with charming penguin for $B \rightarrow PP, PV$ in [22]. The results shown above are obtained with $d_1 = 0.08, d_2 = -1/3$. We see that the branching ratios for $\bar{B}^0 \rightarrow K^- \pi^+, B^- \rightarrow K^- \pi^0$ and $B^- \rightarrow \bar{K}^0 \pi^-$ and the CP asymmetry for $\bar{B}^0 \rightarrow K^- \pi^+$ are in good agreement with experiment. The $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$ is slightly below the measured value. Our results seem reasonable, as the predicted $\mathcal{B}(B^- \rightarrow \pi^- \pi^0)$ which is sensitive $F_0^{B\pi}(0)$ agrees well with the measured value.

Thus QCDF with mainly absorptive additional penguin contribution seems to describe rather well the $B \rightarrow K\pi$ observed branching ratios and the $\bar{B}^0 \rightarrow K^- \pi^+$ direct CP asymmetry. As all the $B \rightarrow K\pi$ branching ratios have been measured with an accuracy at the level of $O(10^{-6})$, one could look at the difference in branching ratios to test QCDF predictions for the electroweak and tree-level contributions. The color-favored electroweak penguin is CKM-favored, so its interference with the strong-penguin terms is about the same size as that of the color-favored tree and penguin interference. For example, from Eq.(5-8), one sees that the differences $2\Gamma(B^- \rightarrow K^- \pi^0) - \Gamma(\bar{B}^0 \rightarrow K^- \pi^+)$ and $2\Gamma(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0) - \Gamma(B^- \rightarrow \bar{K}^0 \pi^-)$ are essentially given by the color-allowed electroweak and strong penguin interference term, so with QCDF expression for the interference term, one could predict $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$ in terms of $\mathcal{B}(B^- \rightarrow \bar{K}^0 \pi^-)$ or vice versa. We have:

$$\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0) = [\mathcal{B}(B^- \rightarrow \bar{K}^0 \pi^-)r_b + \delta B_1]/2 \quad (15)$$

$$\mathcal{B}(B^- \rightarrow K^- \pi^0) = [\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+) + \delta B_2]/2r_b \quad (16)$$

$$\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+) = [\mathcal{B}(B^- \rightarrow \bar{K}^0 \pi^-)r_b + \delta B_3] \quad (17)$$

and where $\delta B_1, \delta B_2, \delta B_3$ are respectively the computed difference $2\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0) - r_b \mathcal{B}(B^- \rightarrow \bar{K}^0 \pi^-)$, $2r_b \mathcal{B}(B^- \rightarrow K^- \pi^0) - \mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+)$, $\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+) - r_b \mathcal{B}(B^- \rightarrow \bar{K}^0 \pi^-)$ and $r_b = \tau_{B^0}/\tau_{B^-}$. We find, in unit of 10^{-6} , with $\rho_A = 1, \phi_A = -55^\circ$ as with




FIG. 1: The computed and measured CP average branching ratios. The horizontal line are the measured values [21] with the gray areas represent the experimental errors. (a), (b), (c), (d) in the left and right figure represent the values for $B^- \rightarrow K^- \pi^0$, $B^- \rightarrow \bar{K}^0 \pi^-$, $\bar{B}^0 \rightarrow K^- \pi^+$ and $\bar{B}^0 \rightarrow \bar{K}^0 \pi^0$ respectively. The curves (a1)-(d1) and (a2)-(d2) are the corresponding QCDF predicted values for $\phi_A = -55^\circ$, without and with additional penguin contribution respectively.




FIG. 2: The same as in Fig.1 but for the computed CP asymmetries.

the computed values in Table I and Table II (only leading terms are shown):

$$\delta B_1 = -3.52 [-3.84(\text{EW}) + 1.42 \cos(\gamma)(\text{C})] \quad (18)$$

$$\delta B_2 = 3.97 [3.84(\text{EW}) - 1.42 \cos(\gamma)(\text{C})] \quad (19)$$

$$\delta B_3 = -1.52 [-5.53 \cos(\gamma)(\text{TP})$$

$$-0.65 \cos(\gamma)(\text{TA}) + 0.69(\text{EWC}) + 0.36(\text{TT})] \quad (20)$$

In the above expressions, the main contributions are from the interference of the color-allowed tree, color-suppressed tree, and electroweak penguin with the strong penguin terms and are shown inside the brackets in the above expressions. Then from Eq.(15-17), we obtain (in unit of 10^{-6})

$$\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0) = 9.0 \pm 0.3, \quad (21)$$

$$\mathcal{B}(B^- \rightarrow K^- \pi^0) = 12.5 \pm 0.3. \quad (22)$$

$$\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+) = 20.0 \pm 0.6, \quad (23)$$

which agrees with experiment, to within the current accuracy. In particular the agreement with experiment

for $\mathcal{B}(B^- \rightarrow K^- \pi^0)$ shows that the electroweak-penguin interference observed in $B \rightarrow K \pi$ decay rates difference is well described by QCDF. The predicted $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$ is a bit lower than experiment, but this prediction is on firm ground since $dB_1 \approx -dB_2$ and the predicted $\mathcal{B}(B^- \rightarrow K^- \pi^0)$ in terms of dB_2 agrees well with experiment, as mentioned above. In fact the discrepancy between experiment and the above prediction for $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$ could be understood in the following way : First of all, Eqs.(18-20) shows that the electroweak-penguin as well as the color-suppressed tree with the strong penguin interference terms in $\mathcal{B}(B^- \rightarrow K^- \pi^0)$ and in $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$ are essentially the same in magnitude but opposite in sign. This can be seen from the decay amplitudes given in Eqs.(5 -8). Then these interference terms cancel out leaving only the color-allowed tree-penguin interference term in the sum of $\Gamma(B^- \rightarrow K^- \pi^0)$ and $\Gamma(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$. Since the color-allowed tree-penguin interference terms, the penguin and annihilation terms are essentially the same in the sum $2(\Gamma(B^- \rightarrow K^- \pi^0) + \Gamma(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0))$ and $(\Gamma(B^- \rightarrow \bar{K}^0 \pi^-) + \Gamma(\bar{B}^0 \rightarrow K^- \pi^+))$, by neglecting small quadratic term in tree and electroweak penguin

contributions, we obtain the approximate relation:

$$2[\mathcal{B}(B^- \rightarrow K^-\pi^0)r_b + \mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)] \\ = [\mathcal{B}(B^- \rightarrow \bar{K}^0\pi^-)r_b + \mathcal{B}(\bar{B}^0 \rightarrow K^-\pi^+)] \quad (24)$$

This relation has been first given in [23, 24], but was derived independently in [25] from the isospin $B \rightarrow K\pi$ amplitudes:

$$\Delta = \{ \Gamma(B^- \rightarrow \bar{K}^0\pi^-) + \Gamma(\bar{B}^0 \rightarrow K^-\pi^+) \\ - 2[\Gamma(B^- \rightarrow K^-\pi^0) + \Gamma(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)] \} \tau_{B^0} \\ = \left[-\frac{4}{3}|B_3|^2 - \frac{8}{\sqrt{3}}\text{Re}(B_3^*B_1e^{i\delta}) \right] (C\tau_{B^0}) \quad (25)$$

where B_1, B_3 are respectively the isospin $I = 1/2$ and $I = 3/2$ tree-level $B \rightarrow K\pi$ amplitudes. With the measured branching ratios, we get $(43.69 \pm 2.4) \times 10^{-6}$ and $(40.9 \pm 2.0) \times 10^{-6}$ for the l.h.s and r.h.s of Eq.(24), and a difference 2.72×10^{-6} between the two values. Ignoring the errors, we see that this difference is rather large, of the size of the tree-penguin interference term, while, theoretically, it should be of $O(10^{-6})$ or less and has a value given by $\delta B_1 + \delta B_2 = 0.43 \times 10^{-6}$. The problem, we suspect, seems to be the measured $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$ which has been lowered over the years. It was $(11.5 \pm 1.0) \times 10^{-6}$ [26] and exceeds the upper limit of $(10.8 \pm 0.3) \times 10^{-6}$ in 2005, as shown in Eq.(15) (the old $K\pi$ puzzle [27]). With our value for $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$, the difference would be reduced to 0.9×10^{-6} and becomes consistent with Eq.(24), within the present accuracy. We thus expect the experimental value for $\mathcal{B}(\bar{B}^0 \rightarrow \bar{K}^0\pi^0)$ would be

lower than the current value and is close to our predicted value $9.0 \pm 0.3 \times 10^{-6}$. (The Belle value is in fact $(9.2 \pm 0.7_{-0.7}^{+0.6}) \times 10^{-6}$ [21]).

III. CONCLUSION

By adding a mainly absorptive, penguin-like contribution to the $B \rightarrow K\pi$ decay QCDF amplitude, we show that QCDF could successfully predict the $B \rightarrow K\pi$ branching ratios. We obtain the correct magnitude and sign for the $\bar{B}^0 \rightarrow K^-\pi^+$ CP asymmetry. The $B^- \rightarrow K^-\pi^0$ CP asymmetry is predicted to be of the same sign and magnitude as that of $\bar{B}^0 \rightarrow K^-\pi^+$ which should be checked against more precise data. We also show that QCDF describes well the electroweak penguin contribution in the $B^- \rightarrow K^-\pi^0$ and $\bar{B}^0 \rightarrow \bar{K}^0\pi^0$ branching ratios and show that the $\bar{B}^0 \rightarrow \bar{K}^0\pi^0$ measured branching ratio should be lower than the current value.

Acknowledgments

I would like to thank Hsiang-nan Li, Hai-Yang Cheng, Chuan-Hung Chen and the members of the Institute of Physics, Academia Sinica, Taiwan, for the warm hospitality and support for my stay at the Institute where part of the work was done. Also thanks to P. Colangelo and F. De Fazio for the warm hospitality and support at INFN, Bari, Italy. This work was supported in part by the EU contract No. MRTN-CT-2006-035482, "FLAVIANet".

-
- [1] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B **606**, 245 (2001).
 - [2] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
 - [3] D. S. Du, J. F. Sun, D. S. Yang and G. H. Zhu, Phys. Rev. D **67**, 014023 (2003);
 - [4] A. Ali, G. Kramer, and C. Lü, Phys. Rev. D **58**, 094009 (1998).
 - [5] This is pointed out by H. Y. Cheng, H. Y. Cheng, private communication.
 - [6] C. Isola, M. Ladisa, G. Nardulli, T. N. Pham, and P. Santorelli, Phys. Rev. D **64** 014029 (2001), *ibid.*, Phys. Rev. D **65** 014005 (2001).
 - [7] T. N. Pham, Proceedings of the QCD 02, Montpellier, Nucl.Phys. (Proc. Suppl.) **B 121** 267 (2003).
 - [8] M. Ciuchini, E. Franco, G. Martinelli, L. Reina, and L. Silvestrini, Phys. Lett. **B 316** 127 (1993); M. Ciuchini, E. Franco, G. Martinelli, and L. Reina, Nucl. Phys. **B 415** 403 (1994).
 - [9] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Lett. **B 597** 291 (2004).
 - [10] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Lett. B **542**, 71 (2002); Phys. Rev. D **69** 054023 (2004).
 - [11] H. Y. Cheng, C. K. Chua and A. Soni, Phys. Rev. D **71**, 014030 (2005).
 - [12] B. Dutta, C. S. Kim, S. Oh, and G. H. Zhu, Eur. Phys. J. C **37**, 273 (2004).
 - [13] D. S. Du, H. Gong, J. F. Sun, D. S. Yang and G. H. Zhu, Phys. Rev. D **65** 074001 (2002).
 - [14] Particle Data Group, *Review of Particle Physics*, Phys. Lett. B **667**, 1 (2008).
 - [15] T.N. Pham, in 2nd Workshop on the CKM Unitarity Triangle, Durham, April 2003, eConf C034052, WG415 (2003), arXiv:hep-ph/0306271.
 - [16] E. Barberio, Talk at the XXIII International Symposium on Lepton and Photon Interactions at High Energy, Aug 13-18 2007, Daegu, Korea.
 - [17] M. Bona *et al.* (UTfit Collaboration), Summary of results at "http://www.utfit.org/".
 - [18] A. Abulencia *et al.*, (CDF Collaboration), Phys. Rev. Lett. **97** 062003 (2006), quoted in Ref.[14].
 - [19] D. Brown, Talk at the XXIII International Symposium on Lepton and Photon Interactions at High Energy, Aug 13-18 2007, Daegu, Korea.
 - [20] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005).
 - [21] Heavy Flavor Averaging Group, E. Barberio *et al.*, arXiv:0808.1297 [hep-ex].
 - [22] N. de Groot, W. N. Cottingham and I. B. Whittingham, Phys. Rev. D **68** 113005 (2003).
 - [23] H. Lipkin, Phys. Lett. **B 445** 403 (1999).

- [24] M. Gronau and J. L. Rosner, Phys. Rev. D **59** 113002 (1999).
- [25] C. Isola and T. N. Pham, Phys. Rev. D **62** 094002 (2000).
- [26] Heavy Flavor Averaging Group, E. Barberio *et al*, hep-ex/0603003.
- [27] S. Baek, C. W. Chiang and D. London, Phys. Lett. **B 675** 59 (2009).